

Simulating the formation of cosmic structure

BY C. S. FRENK

Physics Department, University of Durham, Durham DH1 3LE, England

A timely combination of new theoretical ideas and observational discoveries has brought about significant advances in our understanding of cosmic evolution. Computer simulations have played a key role in these developments by providing the means to interpret astronomical data in the context of physical and cosmological theory. In the current paradigm, our Universe has a flat geometry, is undergoing accelerated expansion and is gravitationally dominated by elementary particles that make up cold dark matter. Within this framework, it is possible to simulate in a computer the emergence of galaxies and other structures from small quantum fluctuations imprinted during an epoch of inflationary expansion shortly after the Big Bang. The simulations must take into account the evolution of the dark matter as well as the gaseous processes involved in the formation of stars and other visible components. Although many unresolved questions remain, a coherent picture for the formation of cosmic structure is now beginning to emerge.

Keywords: Cosmic structure, dark matter, gas dynamics, galaxy formation, computer simulations

1. Introduction

The origin of structure in the Universe is a central problem in Physics. Its solution will not only inform our understanding of the processes by which matter became organized into galaxies and clusters, but it will also help uncover the identity of the dark matter, offer insights into events that happened in the early stages of the Big Bang and provide a useful check on the values of the fundamental cosmological parameters estimated by other means.

Because of its non-linear character, lack of symmetry and general complexity, the formation of cosmic structure is best approached theoretically using numerical simulations. The problem is well posed because the initial conditions – small perturbations in the density and velocity field of matter – are, in principle, known from Big Bang theory and observations of the early Universe, while the basic physical principles involved are understood. The behaviour of the dark matter is governed primarily by gravity, while the formation of the visible parts of galaxies involves gas dynamics and radiative processes of various kinds. Using cosmological simulations it is possible to follow the development of structure from primordial perturbations to the point where the model can be compared with observations.

Over the past few years, there has been huge progress in quantifying observationally the properties of galaxies not only in the nearby universe, but also in the very distant universe. Since the clustering pattern of galaxies is rich with information

about physics and cosmology, much effort is invested in mapping the distribution of galaxies at different epochs. Two large ongoing surveys, the US-based Sloan Digital Sky Survey (York *et al.* 2000), and the Anglo-Australian “2-degree field galaxy redshift survey” (2dFGRS, Colless *et al.* 2001), are revolutionizing our view of the nearby universe with order of magnitude increases in the amount of available data. Similarly, new data collected in the past five years or so have, for the first time, opened up the high redshift universe† to detailed statistical study (Steidel *et al.* 1996).

The advent of large computers, particularly parallel supercomputers, together with the development of efficient algorithms, has enabled the accuracy and realism of simulations to keep pace with observational progress. With the wealth of data now available, simulations are essential to interpret astronomical data and to link them to physical and cosmological theory.

2. Building a model

To build a model of large-scale structure, four key ingredients need to be specified: (i) the content of Universe, (ii) the initial conditions, (iii) the growth mechanism, and (iv) the values of fundamental cosmological parameters. I now discuss each of these in turn.

(a) *The content of the Universe*

Densities are usually expressed in terms of the cosmological density parameter, $\Omega = \rho/\rho_{crit}$, where the critical density, ρ_{crit} , is the value that makes the geometry of the Universe flat. The main constituents of the Universe and their contribution to Ω are listed in Table 1.

Table 1. *The content of the Universe*

Component	Contribution to Ω
CMB radiation	$\Omega_r = 4.7 \times 10^{-5}$
massless neutrinos	$\Omega_\nu = 3 \times 10^{-5}$
massive neutrinos	$\Omega_\nu = 6 \times 10^{-2} \left(\frac{\langle m_\nu \rangle}{1\text{ev}} \right)$
baryons	$\Omega_b = 0.037 \pm 0.009$
(of which stars)	$\Omega_s = (0.0023 - 0.0041) \pm 0.0004$
dark matter	$\Omega_{dm} \simeq 0.3$
dark energy	$\Omega_\Lambda \simeq 0.7$

The main contribution to the extragalactic radiation field today is the cosmic microwave background (CMB), the redshifted radiation left over from the Big Bang. These photons have been propagating freely since the epoch of “recombination”, approximately 300,000 years after the Big Bang. The CMB provides a direct observational window to the conditions that prevailed in the early Universe. The Big

† In cosmology, distances to galaxies are estimated from the redshift of their spectral lines; higher redshifts correspond to more distant galaxies and thus to earlier epochs.

Bang also produced neutrinos which today have an abundance comparable to that of photons. We do not yet know for certain what, if any, is the mass of the neutrino, but even for the largest masses that seem plausible at present, $\sim 0.1\text{eV}$, neutrinos make a negligible contribution to the total mass budget (although they could be as important as baryons). The abundance of baryons is now known with reasonable precision from comparing the abundance of deuterium predicted by Big Bang theory with observations of the absorption lines produced by intergalactic gas clouds at high redshift seen along the line-of-sight to quasars (Tytler *et al.* 2000). Baryons, the overwhelming majority of which are *not* in stars today, are also dynamically unimportant (except, perhaps, in the cores of galaxies).

Dark matter makes up most of the matter content of the Universe today. To the now firm dynamical evidence for its existence in galaxy halos, even more direct evidence has been added by the phenomenon of gravitational lensing which has now been detected around galaxy halos (e.g. Fischer *et al.* 2000, McKay *et al.* 2001, Wilson *et al.* 2001), in galaxy clusters (e.g. Clowe *et al.* 2000), and in the general mass field (e.g. Van Waerbeke *et al.* 2001 and references therein). The distribution of dark matter in rich clusters can be reconstructed in fair detail from the weak lensing of distant background galaxies in what amounts virtually to imaging the cluster dark matter. Various dynamical tests are converging on a value of $\Omega_{\text{dm}} \simeq 0.3$, which is also consistent with independent determinations such as those based on the baryon fraction in clusters (White *et al.* 1993, Evrard 1997), and on the evolution in the abundance of galaxy clusters (Eke *et al.* 1998, Borgani *et al.* 2001). Since Ω_{dm} is much larger than Ω_b , it follows that the dark matter cannot be made of baryons. The most popular candidate for the dark matter is a hypothetical elementary particle like those predicted by supersymmetric theories of particle physics. These particles are referred to generically as cold dark matter or CDM. (Hot dark matter is also possible, for example, if the neutrino had a mass of ~ 5 eV. However, early cosmological simulations showed that the galaxy distribution in a universe dominated by hot dark matter would not resemble that observed in our Universe (White, Frenk and Davis 1983).)

A recent addition to the cosmic budget is the dark energy, direct evidence for which was first provided by studies of type Ia supernovae (Riess *et al.* 1998, Perlmutter *et al.* 1999)†. These presumed ‘standard candles’ can now be observed at redshifts between 0.5 and 1 and beyond. The more distant ones are fainter than would be expected if the universal expansion were decelerating today, indicating that the expansion is, in fact, accelerating. Within the standard Friedmann cosmology, there is only one agent that can produce an accelerating expansion. This is nowadays known as dark energy, a generalization of the cosmological constant first introduced by Einstein, which could, in principle, vary with time. The supernova evidence is consistent with the value $\Omega_\Lambda \simeq 0.7$. Further, independent evidence for dark energy is provided by a recent joint analysis of CMB data (see next section) and the 2dFGRS (Efstathiou *et al.* 2002).

Amazingly, when all the components are added together, the data are consistent with a flat universe:

† The possibility that dark energy might be the dynamically dominant component had been anticipated by theorists from studies of the cosmic large-scale structure (see e.g. Efstathiou *et al.* 1990), and was considered in the first simulations of structure formation in cold dark matter universes (Davis *et al.* 1985).

$$\Omega = \Omega_b + \Omega_{\text{dm}} + \Omega_\Lambda \simeq 1 \quad (2.1)$$

(b) *The initial conditions*

The idea that galaxies and other cosmic structures are the result of the slow amplification by the force of gravity of small primordial perturbations present in the mass density at early times goes back, at least, to the 1940s (Lifshitz 1946). However, it was only in the early 1980s that a physical mechanism capable of producing small perturbations was identified. This is the mechanism of inflation, an idea due to Guth (1981), which changed the face of modern cosmology. Inflation is produced by the dominant presence of a quantum scalar field which rolls slowly from a false to the true vacuum, maintaining its energy density approximately constant and causing the early Universe to expand exponentially for a brief period of time. Quantum fluctuations in the inflaton field are blown up to macroscopic scales and become established as genuine adiabatic ripples in the energy density. Simple models of inflation predict the general properties of the resulting fluctuation field: it has Gaussian distributed amplitudes and a near scale-invariant power spectrum (Starobinskii 1982).

After three decades of ever more sensitive searches, evidence for the presence of small fluctuations in the early universe was finally obtained in 1992. Since prior to recombination the matter and radiation fields were coupled, fluctuations in the mass density are reflected in the temperature of the radiation. Temperature fluctuations in the CMB were discovered by the COBE satellite (Smoot *et al.* 1992) and are now being measured with ever increasing accuracy, particularly by detectors deployed in long-flight balloons (de Bernardis *et al.* 2000, Hanany *et al.* 2000, Leitch *et al.* 2002). The spectrum of temperature fluctuations is just what inflation predicts: it is scale invariant on large scales and shows a series of ‘‘Doppler’’ or ‘‘acoustic’’ peaks which are the result of coherent acoustic oscillations experienced by the photon-baryon fluid before recombination. The characteristics of these peaks depend on the values of the cosmological parameters. For example, the location of the first peak is primarily determined by the large-scale geometry of the Universe and thus by the value of Ω . Current data imply a flat geometry, consistent with eqn. 2.1.

The spectrum of primordial fluctuations generated, for example, by inflation evolves with time in a manner that depends on the content of the Universe and the values of the cosmological parameters. The dark matter acts as a sort of filter, inhibiting the growth of certain wavelengths and promoting the growth of others. Following the classical work of Bardeen *et al.* (1986), transfer functions for different kinds of dark matter (and different types of primordial fluctuation fields, including non-Gaussian cases) have been computed. In Gaussian models, the product (in Fourier space) of the primordial spectrum and the transfer function, together with the growing mode of the associated velocity field, provides the initial conditions for the formation of cosmic structure.

(c) *Growth mechanism*

Primordial fluctuations grow by gravitational instability: overdense fluctuations expand linearly, at a retarded rate relative to the Universe as a whole, until even-

tually they reach a maximum size and collapse non-linearly to form an equilibrium (or ‘virialized’) object whose radius is approximately half the physical size of the perturbation at maximum expansion. The theory of fluctuation growth is lucidly explained by Peebles (1980).

Although gravitational instability is now widely accepted as the primary growth mechanism responsible for the formation of structure, it is only very recently that firm empirical evidence for this process was found. Gravitational instability causes inflow of material around overdense regions. From the perspective of a distant observer, this flow gives rise to a characteristic infall pattern which is, in principle, measurable in a galaxy redshift survey by comparing the two-point galaxy correlation function along and perpendicular to the line-of-sight. In this space, the infall pattern resembles a butterfly (Kaiser 1987). This pattern has been clearly seen for the first time in the 2dFGRS (Peacock *et al.* 2001)†.

(d) Cosmological parameters

After decades of debate, the values of the fundamental cosmological parameters are finally being measured with some degree of precision. The main reason for this is the accurate measurement of the acoustic peaks in the CMB temperature anisotropy spectrum whose location, height and shape depend on the values of the cosmological parameters. Some parameter degeneracies exist but some of these can be broken using other data, for example, the distant Type Ia supernovae or the 2dFGRS (eg. Efstathiou *et al.* 2002). The CMB data alone do not constrain the Hubble constant, but there is a growing consensus from the HST key project (Freedman *et al.* 2001), and other methods, that its value, in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is $h = 0.70 \pm 0.07$. In addition to h and the other parameters listed in Table 1, the other important number in studies of large-scale structure is the amplitude of primordial density fluctuations which is usually parametrized by the quantity σ_8 (the linearly extrapolated value of the top-hat filtered fluctuation amplitude on the fiducial scale of $8 h^{-1} \text{ Mpc}$). The best estimate of this quantity comes from the observed abundance of rich galaxy clusters which gives $\sigma_8 \Omega^{0.6} = 0.5$, with an uncertainty of about 10% (Eke, Cole & Frenk 1996, Viana & Liddle 1996, Pierpaoli *et al.* 2001).

3. Cosmological simulations

Operationally, the problem of the cosmic large-scale structure can be divided into two parts: understanding the clustering evolution of the dark matter and understanding the gaseous and radiative processes that lead to the formation of galaxies. Specialized simulation techniques have been developed to tackle both aspects of the problem. The evolution of the dark matter is most often calculated using N-body techniques, implemented through a variety of efficient algorithms, such as P³M (Particle-particle/particle-mesh; Efstathiou *et al.* 1985), AP³M (the adaptive mesh version of P³M; Couchman *et al.* 1995) and hierarchical trees (Barnes & Hut 1986, Springel *et al.* 2001, Stadel 2000). Gaseous and radiative processes

† Strictly speaking the ‘butterfly’ pattern does not prove the existence of infall since the continuity equation would ensure a similar pattern even if velocities were induced by non-gravitational processes. However, it can be shown that such velocities, if present, would rapidly decay.

are followed by combining a hydrodynamics code with an N-body code. Numerical hydrodynamic techniques used in cosmology include Eulerian methods (Cen 1992), Lagrangian codes based on Smooth Particle Hydrodynamics (SPH) (Gingold & Monaghan 1977), and hybrid codes (e.g. Gnedin 1995, Pen 1998). These techniques have different strengths and weaknesses, but they all give similar results in the simplest cosmological problems where a detailed comparison has been performed (Frenk *et al.* 1999).

There has been a rapid growth in the size and power of cosmological simulations in the two and a half decades since this technique was introduced into the subject by Peebles (1970). One way to measure this growth is by the number of particles employed in the simulations. The size of the largest simulations has grown exponentially, in a manner reminiscent of the well-known “Moore’s law” that describes the increase in cpu speed with time, except that the advent of massively parallel supercomputers led to a sudden order-of-magnitude jump in size towards the end of the past decade. The largest simulations carried out to date are the 1-billion particle “Hubble volume,” N-body simulations performed by the Virgo consortium, an international collaboration of researchers in the UK, Germany and Canada.

(a) *Large-scale structure*

Figure 1 illustrates the spatial distribution of *dark matter* at the present day, in a series of simulations covering a large range of scales. Each panel is a thin slice of the cubical simulation volume and shows the slightly smoothed density field defined by the dark matter particles. In all cases, the simulations pertain to the “ Λ CDM” cosmology, a flat cold dark matter model in which $\Omega_{\text{dm}} = 0.3$, $\Omega_{\Lambda} = 0.7$ and $h = 0.7$. The top-left panel illustrates the Hubble volume simulation: on these large scales, the distribution is very smooth. To reveal more interesting structure, the top right panel displays the dark matter distribution in a slice from a volume approximately 2000 times smaller. At this resolution, the characteristic filamentary appearance of the dark matter distribution is clearly visible. In the bottom-right panel, we zoom again, this time by a factor of 5.7 in volume. We can now see individual galactic-size halos which preferentially occur along the filaments, at the intersection of which large halos form that will host galaxy clusters. Finally, the bottom-left panel zooms into an individual galactic-size halo. This shows a large number of small substructures that survive the collapse of the halo and make up about 10% of the total mass (Klypin *et al.* 1999, Moore *et al.* 1999)

For simulations like the ones illustrated in Figure 1, it is possible to characterize the statistical properties of the dark matter distribution with very high accuracy. For example, Figure 2 shows the 2-point correlation function, $\xi(r)$, of the dark matter (a measure of its clustering strength) in the simulation depicted in the top-right of Figure 1 (Jenkins *et al.* 1998). The statistical error bars in this estimate are actually smaller than the thickness of the line. Similarly, higher order clustering statistics, topological measures, the mass function and clustering of dark matter halos and the time evolution of these quantities can all be determined very precisely from these simulations (e.g. Jenkins *et al.* 2001, Evrard *et al.* 2002). In a sense, the




Figure 1. Slices through 4 different simulations of the dark matter in the ‘ Λ CDM’ cosmology. Denoting the number of particles in each simulation by N , the length of the simulation cube by L , the thickness of the slice by t , and the particle mass by m_p , the characteristics of each panel are as follows. Top-left (the Hubble volume simulation, Evrard *et al.* 2002): $N = 10^9$, $L = 3000 h^{-1}$ Mpc, $t = 30 h^{-1}$ Mpc, $m_p = 2.2 \times 10^{12} h^{-1} M_\odot$. Top-right (Jenkins *et al.* 1998): $N = 16.8 \times 10^6$, $L = 250 h^{-1}$ Mpc, $t = 25 h^{-1}$ Mpc, $m_p = 6.9 \times 10^{10} h^{-1} M_\odot$. Bottom-right (Jenkins *et al.* 1998): $N = 16.8 \times 10^6$, $L = 140 h^{-1}$ Mpc, $t = 14 h^{-1}$ Mpc, $m_p = 1.4 \times 10^{10} h^{-1} M_\odot$. Bottom-left (Navarro *et al.* 2002): $N = 7 \times 10^6$, $L = 0.5 h^{-1}$ Mpc, $t = 1 h^{-1}$ Mpc, $m_p = 6.5 \times 10^5 h^{-1} M_\odot$.

problem of the distribution of dark matter in the Λ CDM model can be regarded as largely solved[†].

[†] However, the innermost structure of halos like those in the bottom-left of Figure 1 is still a matter of controversy.




Figure 2. Two-point correlation functions. The dotted line shows the dark matter $\xi_{dm}(r)$ (Jenkins *et al.* 1998). The solid line shows the galaxy predictions of Benson *et al.* (2000), with Poisson errors indicated by the dashed lines. The points with errorbars show the observed galaxy $\xi_{gal}(r)$ (Baugh 1996). The galaxy data are discussed in §3(b). (Adapted from Benson *et al.* (2001a).

In contrast to the clustering of the dark matter, the process of galaxy formation is still poorly understood. How then can dark matter simulations like those of Figure 1 be compared with observational data which, for the most part, refer to galaxies? On large scales a very important simplification applies: for Gaussian theories like CDM, it can be shown that if galaxy formation is a local process, that is, if it depends only upon local physical conditions (density, temperature, etc), then, on scales much larger than that associated with individual galaxies, the galaxies must trace the mass, i.e. on sufficiently large scales, $\xi_{gal}(r) \propto \xi_{dm}(r)$ (Coles 1993). It suffices therefore to identify a random subset of the dark matter particles in the simulation to obtain an accurate prediction for the properties of galaxy clustering on large scales. This idea (complemented on small scales by an empirical prescription in the manner described by Cole *et al.* 1998) has been used to construct the mock versions of a region of the APM galaxy survey and of a slice of the 2dFGRS displayed in Figures 3 and 4 which also show the real data for comparison in each case. By eye at least, it is very difficult to distinguish the mocks from the real data.

A quantitative comparison between simulations and the real world is carried out in Figure 5. The symbols show the estimate of the power spectrum in the 2dFGRS survey (Percival *et al.* 2001). This is the raw power spectrum convolved with the survey window function and can be compared directly with the line showing the theoretical prediction obtained from the mock catalogues which have exactly the same window function. The agreement between the data and the Λ CDM model is remarkably good.




Figure 3. The region of the APM projected galaxy survey from which the 2dFGRS is drawn. Only galaxies brighter than $m_{b_j} = 19.35$ are plotted. The top panel is the real data and the other two panels are mock catalogues constructed from the Hubble volume simulations.

(b) Galaxy formation

Understanding galaxy formation is a much more difficult problem than understanding the evolution of the dark matter distribution. In the CDM theory, galaxies form when gas, initially well mixed with the dark matter, cools and condenses into emerging dark matter halos. In addition to gravity, a non-exhaustive list of the processes that now need to be taken into account includes: the shock heating and cooling of gas into dark halos, the formation of stars from cold gas and the evolution of the resulting stellar population, the feedback processes generated by the ejection of mass and energy from evolving stars, the production and mixing of heavy elements, the extinction and reradiation of stellar light by dust particles, the formation of black holes at the centres of galaxies and the influence of the associated quasar emission. These processes span an enormous range of length and mass scales. For




Figure 4. A 1° thick slice through the 2dF galaxy redshift survey. The radial coordinate is redshift and the angular coordinate is right ascension. The top-left panel is the real data and the other two panels are mock catalogues constructed from the Hubble volume simulations.

example, the parsec scale relevant to star formation is a factor of 10^8 smaller than the scale of a galaxy supercluster.

The best that can be done with current computing techniques is to model the evolution of dark matter and gas in a cosmological volume with resolution comparable to a single galaxy. Subgalactic scales must then be regarded as “sub-grid” scales and followed by means of phenomenological models based either on our current physical understanding or on observations. In the approach known as “semi-analytic” modelling (White & Frenk 1991), even the gas dynamics is treated phenomenologically using a simple, spherically symmetric model to describe the accretion and cooling of gas into dark matter halos. It turns out that this simple model works surprisingly well as judged by the good agreement with results of full N-body/gas-dynamical simulations (Benson *et al.* 2001b, Helly *et al.* 2002, Yoshida *et al.* 2002).




Figure 5. The power spectrum of the 2dFGRS (symbols) compared with the power spectrum predicted in the Λ CDM model (line). Both power spectra are convolved with the 2dFGRS window function. The model predictions come from dark matter simulations and assume that, on large scales, the distribution of galaxies traces the distribution of mass. (Adapted from Percival *et al.* 2001).

The main difficulty encountered in cosmological gas dynamical simulations arises from the need to suppress a cooling instability present in hierarchical clustering models like CDM. The building blocks of galaxies are small clumps that condense at early times. The gas that cools within them has very high density, reflecting the mean density of the Universe at that epoch. Since the cooling rate is proportional to the square of the gas density, in the absence of heat sources, most of the gas would cool in the highest levels of the mass hierarchy leaving no gas to power star formation today or even to provide the hot, X-ray emitting plasma detected in galaxy clusters. Known heat sources are photoionisation by early generations of stars and quasars and the injection of energy from supernovae and active galactic nuclei. These processes, which undoubtedly happened in our Universe, belong to the realm of subgrid physics which cosmological simulations cannot resolve. Different treatments of this “feedback” result in different amounts of cool gas and can lead to very different predictions for the properties of the galaxy population. This is a fundamental problem that afflicts cosmological simulations even when they are complemented by the inclusion of semi-analytic techniques. In this case, the resolution of the calculation can be extended to arbitrarily small mass halos, perhaps allowing a more realistic treatment of feedback. Although they are less general than full gasdynamical simulations, simulations in which the evolution of gas is treated semi-analytically make experimentation with different prescriptions relatively simple and efficient (Kauffmann White & Guiderdoni 1993, Somerville & Primack 1999, Cole *et al.* 2000)

The outcome of an N-body dark matter simulation in a Λ CDM universe in which the visible properties of the galaxies have been calculated using the semi-analytic model of Cole *et al.* (2000) is illustrated in Fig. 6 (Benson *et al.* 2001a). Galaxies




Figure 6. A slice $10 h^{-1}$ Mpc thick of a simulation of a cubic region of side $141 h^{-1}$ Mpc in the Λ CDM cosmology. The grey scale shows a slightly smoothed representation of the dark matter in the N-body simulation. The coloured dots show galaxies; the size of the dots is proportional to the B-band luminosity of the galaxy and the colour represents the B-V colour as given on the scale on the top. The top panel corresponds to redshift $z = 0$ and the bottom panel to $z = 3$. (Adapted from Benson *et al.* 2001a).

form mostly along the filaments delineated by the dark matter. Red galaxies predominate in the most massive dark matter halos, just as observed in real galaxy clusters. This segregation is a natural outcome of hierarchical clustering from CDM initial conditions. It reflects the fact that the progenitors of rich clusters form substantially earlier than a typical dark matter halo of the same mass. Fig. 7 shows the galaxy luminosity function which describes the abundance of galaxies of different luminosities. The theoretical predictions, shown by the line, agree remarkably well with the observations but this should not be regarded as a spectacular success of the theory because the free parameters in the semi-analytic star formation and feedback model have been tuned to achieve as good a match as possible to this




Figure 7. The galaxy luminosity function. The symbols show the number of galaxies per unit volume and per unit magnitude measured in various surveys, as a function of galaxy magnitude (open circles: Zucca *et al.* 1997; open squares: Loveday *et al.* 1992; thick error bars: Norberg *et al.* 2001b). The solid line shows the predictions of the semi-analytic model of Cole *et al.* (2000).

specific observational dataset. In particular, the feedback model has been tuned to produce a relatively flat function at the faint end.

Having fixed the model parameters by reference to a small subset of the data such as the galaxy luminosity function, we can ask whether the same model accounts for other basic observational data. The galaxy autocorrelation function, $\xi_{gal}(r)$, in the simulations is plotted in Fig. 2 above. On large scales, it follows $\xi_{dm}(r)$ quite closely, but on small scales it dips below the mass autocorrelation function. This small scale “antibias” has also been seen in N-body/gasdynamical simulations of the Λ CDM cosmology (Pearce *et al.* 1999, 2001, Dave *et al.* 1999), and in dark matter simulations that resolve individual galactic halos (Klypin *et al.* 1999). The galaxy autocorrelation function in the simulations of Benson *et al.* (2000) agrees remarkably well with the observational data (see also Kauffmann *et al.* 1999a). This is a genuine success of the theory because no model parameters have been adjusted in this comparison. The differences between the small-scale clustering of galaxies and dark matter result from the interplay between the clustering of dark matter halos and the occupation statistics of galaxies in halos which, in turn, are determined by the physics of galaxy formation. This conclusion, discussed in detail by Benson *et al.* (2000), has led to the development of an analytic formulation known as the “halo model” (e.g. Seljak 2000, Peacock & Smith 2000, Berlind & Weinberg 2002).

Another genuine prediction of the model is the dependence of the strength of clustering on the luminosity of different subsamples. It can be seen in Fig. 6 that




Figure 8. The correlation length as a function of the luminosity of different galaxy subsamples. The correlation length is defined as the pair separation for which $\xi(r) = 1$. The symbols show the results from the 2dFGRS and the line the predictions of the simulations of Benson *et al.* (2000). (Adapted from Norberg *et al.* 2001a).

the brightest galaxies are concentrated in the most massive clusters, leading one to suspect that their autocorrelation function must be stronger than average. This is indeed the case, as illustrated in Fig. 8 which compares the variation of the clustering length (defined as the pair separation for which $\xi(r) = 1$) of galaxy samples of different intrinsic luminosity in the simulations of Benson *et al.* (2001a) with the observational data obtained from the 2dFGRS by Norberg *et al.* (2001a). The agreement between theory and observations is remarkable considering that there are no adjustable parameters in this comparison. The reason for the strong clustering of bright galaxies is related to the colour-density relation seen in Fig. 6: the brightest galaxies form in the highest peaks of the density distribution which, in initially Gaussian fields, are more strongly clustered than average peaks which produce less extreme galaxies.

The patch of model universe illustrated in the top panel of Fig. 6 is shown at the earlier epoch corresponding to redshift $z = 3$ (when the universe was only about 20% of its current age) in the bottom panel of this figure. The galaxies are now blue, reflecting the colour of their younger stellar population. There are fewer galaxies in this plot than in the $z = 0$ slice. In fact, this is the epoch when the first substantial population of bright galaxies formed in the simulation. As Baugh *et al.* (1998) argued, the properties of these model galaxies resemble those of the “Lyman-break” galaxies discovered by Steidel *et al.* (1996), even though different models make somewhat different predictions for their exact properties (Somerville *et al.* 2001). Most models, however, predict that the brightest galaxies at $z = 3$ should be strongly clustered (Kauffmann *et al.* 1999b) and, indeed, the models of Baugh *et al.* (1998) correctly anticipated that the Lyman-break galaxies would

have a clustering length comparable to that of bright galaxies today (Adelberger *et al.* 1998). This too should be regarded as a significant success of this kind of modelling in the Λ CDM cosmology. As Fig. 6 shows, in contrast to the galaxies, the dark matter is much more weakly clustered at $z = 3$ than at $z = 0$, indicating that galaxies were strongly biased at birth.

4. Conclusions

Unlike most computational problems in many areas of science, the cosmological problem is blessed with known, well-specified initial conditions. Within a general class of models, it is possible to calculate the properties of primordial perturbations in the cosmic energy density generated by quantum processes during an early inflationary epoch. In a wide family of inflationary models, these perturbations are adiabatic, scale-invariant and have Gaussian-distributed Fourier amplitudes. The model also requires an assumption about the nature of the dark matter and the possibilities have now been narrowed down to non-baryonic candidates of which cold dark matter particles seem the most promising. An empirical test of the initial conditions for the formation of structure predicted by the model is provided by the cosmic microwave background radiation. The tiny temperature fluctuations it exhibits have exactly the properties expected in the model. Furthermore, the CMB data can be used to fix some of the key model parameters such as Ω and Ω_b , while these data, combined with other recent datasets such as the 2dFGRS, allow the determination of many of the remaining parameters such as Ω_m , Ω_Λ and h . It is this specificity of the cosmological problem that has turned simulations into the primary tool for connecting cosmological theory to astronomical observations.

In addition to well-specified initial conditions, the cosmological dark matter problem has the advantage that the only physical interaction that is important is gravity. The problem can thus be posed as a gravitational N-body problem and approached using the many sophisticated techniques that have been developed over the past two decades to tackle this problem. Although on small scales there remain a number of unresolved issues, it is fair to say that on scales larger than a few megaparsecs, the distribution of dark matter in CDM models is essentially understood. The inner structure of dark matter halos, on the other hand, is still a matter of debate and the mass function of dark matter halos has only been reliably established by simulations down to masses of order $10^{11} M_\odot$. Resolving these outstanding issues is certainly within reach, but this will require carefully designed simulations and large amounts of computing power.

The frontier of the subject at present lies in simulations of the formation, evolution and structure of galaxies. This problem requires first of all a treatment of gas dynamics in a cosmological context and a number of techniques, relying on direct simulations or on semi-analytical approximations, are being explored. There are quite a few different approaches to cosmological gasdynamics, but it is reassuring that they all give similar results in the simplest relevant problem, the evolution of non-radiative gas during the formation of a galaxy cluster. No detailed comparisons exist yet for the more complicated case in which the gas is allowed to cool, but at least one of the gasdynamic simulation techniques, SPH, gives quite similar results to a simple semi-analytic approach. Realistic models of galaxy formation, however, will require much more than a correct treatment of cooling gas. Such mod-

els will necessarily have to include a plethora of astrophysical phenomena such as star formation, feedback, metal enrichment, etc. The huge disparity between the submegaparsec scales on which these processes operate and the gigaparsec scale of the large-scale structure makes it impossible to contemplate a comprehensive *ab initio* calculation. The way forward is clearly through a hybrid approach combining direct simulation of processes operating on a limited range of scales with a phenomenological treatment of the others. There is currently a great deal of activity in the phenomenology of galaxy formation.

In spite of the uncertainties that remain, all the indications are that our Universe is well described by a model in which

- (i) the overall geometry is flat;
- (ii) the dominant dynamical components are cold dark matter ($\sim 30\%$) and dark energy ($\sim 70\%$) with baryons playing very much a supporting role ($\sim 4\%$);
- (iii) the initial conditions are quantum fluctuations in the primordial energy density generated during inflation and
- (iv) structure has grown primarily as a result of the gravitational instability experienced by mass fluctuations in an expanding universe.

A skeptic is entitled to feel that the current paradigm is odd, to say the least. Not only is there a need to invoke vast amounts of as yet undetected non-baryonic cold dark matter, but there is also the need to account for the dominant presence of a dark energy whose very existence is a mystery within conventional models of fundamental physics. Odd as it may seem, however, this model accounts remarkably well for a large and diverse collection of empirical facts that span 13 billion years of evolution.

5. Acknowledgements

I am grateful to my collaborators for their contribution to the work reviewed here, especially Carlton Baugh, Andrew Benson, Shaun Cole, Adrian Jenkins, Cedric Lacey, Peder Norberg, John Peacock, Will Percival, and Simon White.

6. References

- Adelberger K. L., Steidel C. C., Giavalisco M., Dickinson M., Pettini M., Kellogg M. (1998), *ApJ*, 505, 18
- Bardeen, J. M., Bond, J. R., Kaiser, N. and Szalay, A. S. (1986), *Ap.J.*, 304, 15
- Barnes, J., Hut, P. (1986), *Nature*, 324, 446
- Baugh C. M. (1996), *MNRAS*, 280, 26
- Baugh C. M., Cole S., Frenk C. S. (1996b), *MNRAS*, 283, 1361
- Baugh C. M., Cole S., Frenk C. S. Lacey C. G., (1998), *ApJ*, 498, 504
- Benson A. J., Cole S., Frenk C. S., Baugh C. M., Lacey C. G. (2000), *MNRAS*, 311, 79
- Benson, A. J., Frenk, C. S., Baugh, C. M., Cole, S., Lacey, C. G. (2001a), *MNRAS*, 327, 1041

- Benson, A. J., Pearce, F. R., Frenk, C. S., Baugh, C. M. & Jenkins, A. R. (2001b), MNRAS, 320, 261
- Berlind, A. & Weinberg, D. (2002), ApJ, in press (astro-ph/0109001)
- de Bernardis, P. et al (2000), Nature 404, 995
- Borgani, S. et al (2001), ApJ, 561, 13
- Cen, R. (1992), ApJ Suppl 78, 341
- Clowe, D., Luppino, G. A., Kaiser, N., Gioia, I. M. (2000), ApJ, 539, 540
- Cole, S., Hatton, S., Weinberg, D. H., Frenk, C. S. (1998), MNRAS, 300, 945
- Cole, S., Lacey, C. G., Baugh, C. M., Frenk, C.S. (2000), MNRAS, 319, 168
- Coles, P. (1993) MNRAS, 262, 1065
- Colless, M. et al (the 2dFGRS team) (2001), MNRAS, 328, 1039
- Couchman, H. M. P., Thomas, P. A., Pearce, F. R. (1995), ApJ, 452, 797
- Dave, R., Hernquist, L., Katz, N. & Weinberg, D. (1999), in Proceedings of Rencontres Internationales de l'IGRAP, Clustering at High Redshift, Marseille 1999, astro-ph/9910221
- Davis, M., Efstathiou, G., Frenk, C. S. & White, S. D. M. (1985), ApJ., 292, 371
- Efstathiou, G., Davis, M., Frenk, C. S. & White, S. D. M. (1985), ApJ Suppl, 57, 241
- Efstathiou, G. et al (The 2dFGRS team) (2002), MNRAS, in press, (astro-ph/0109152)
- Efstathiou, G., Sutherland, W. J., Maddox, S. J. (1990), Nature, 348, 705
- Eke, V. R., Cole, S. & Frenk, C. S. (1996), MNRAS, 282, 263
- Eke, V. R., Cole, S. Frenk, C. S. & Henry, J. P. (1998), MNRAS, 298, 1145
- Evrard, A. E. (1997), MNRAS, 292, 289.
- Evrard, A.E., MacFarland, T., Couchman, H.M.P., Colberg, J.M., Yoshida, N., White, S.D.M., Jenkins, A., Frenk, C.S., Pearce, F.R., Efstathiou, G., Peacock, J. & Thomas, P., (The Virgo Consortium) (2002), ApJ, in press.
- Fischer, P. et al (2000), AJ 120, 1198
- Frenk, C. S. et al (1999), ApJ, 525, 554
- Freedman, W. L. et al (2001), ApJ, 553, 47
- Gingold, R. A. & Monaghan, J. J. (1977), MNRAS 181, 375
- Gnedin, N. Y. (1995), ApJ Suppl, 97, 231
- Guth, A. H. (1981), Phys Rev D, 23, 347
- Hanany, S. et al (2000), ApJ Lett, 545, 5
- Helly, J., Cole, S. M., Frenk, C. S., Baugh, C., Benson, A. & Lacey, C. G. (2002), submitted to MNRAS (astro-ph/0202485)
- Jenkins, A., Frenk, C. S., Pearce, F. R., Thomas, P., Colberg, J., White, S. D. M., Couchman, H., Peacock, J., Efstathiou, G. & Nelson, A. (1998), ApJ, 499, 20
- Jenkins A., Frenk C. S., White S. D. M., Colberg J. M., Cole S., Evrard A. E., Yoshida N. (2001), MNRAS, 321,372
- Kaiser, N. (1987), MNRAS, 227, 1
- Kauffmann, G., Colberg, J., Diaferio, A., White, S. D. M. (1999a), MNRAS, 303, 188

- Kauffmann, G., Colberg, J., Diaferio, A., White, S. D. M. (1999b), MNRAS, 307, 529
- Kauffmann G., White S. D. M., Guiderdoni B. (1993), MNRAS, 264, 201
- Klypin, A., Gottlober, S., Kravstov, A., Khokhlov, A.M. (1999), ApJ, 516, 530
- Leitch, E.M. et al (2002), ApJ, 568, 28L
- Lifshitz, E. M. (1946), J Phys, 10, 116
- Loveday, J., Peterson, B., Efstathiou, G. & Maddox, S. (1992), ApJ, 319, 338
- McKay, T. et al (2001) ApJ, in press (astro-ph/0108013)
- Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., Tozzi, P. (1999), ApJ, 524, 19
- Navarro, J. F., Power, C., Jenkins, A. R., Frenk, C. S. & White, S. D. M. (2002), in preparation
- Norberg, P., Baugh, C. M., Hawkins, E., Maddox, S., Peacock, J. Cole, S., Frenk, C. S., et al (The 2dFGRS team) (2001a), MNRAS, 328, 64
- Norberg, P., Cole, S., Baugh, C. M., Frenk, C. S. et al (The 2dFGRS team) (2001b), submitted to MNRAS, (astro-ph/0111011)
- Peacock, J. A. & Smith, R. (2000) 318, 1144
- Peacock, J. et al (The 2dFGRS team) (2001), Nature, 410, 169
- Pearce, F. R., Jenkins, A. R., Frenk, C. S., Colberg, J., White, S. D. M., Thomas, P., Couchman, H. M. P., Peacock, J. A., Efstathiou, G. (the Virgo Consortium) (1999), ApJ, 521, 992
- Pearce, F. R., Jenkins, A. R., Frenk, C. S., White, S. D. M., Thomas, P., Couchman, H. M. P., Peacock, J. A., Efstathiou, G. (the Virgo Consortium) (2001), MNRAS, 326, 649
- Peebles, P. J. E. (1970), AJ, 75, 13
- Peebles, P. J. E. (1980), The large scale structure of the Universe. (Princeton: Princeton University Press).
- Pen, U-L. (1998), ApJ Suppl, 115, 19
- Percival, W. et al (the 2dFGRS team) 2001, MNRAS, 327, 1297
- Perlmutter, S. et al (The Supernova Cosmology Project) (1999), ApJ. 517, 565
- Pierpaoli, E., Scott, D. & White, M. (2001), MNRAS 325, 77
- Riess, A. et al (1998), AJ, 116, 1009
- Seljak, U. (2000), MNRAS, 318, 203
- Smoot, G. et al (1992), Apj 396, 1
- Somerville R. S., Primack J. R. (1999), MNRAS, 310, 1087
- Somerville, Rachel S., Primack, Joel R., Faber, S. M. (2001), MNRAS, 320, 504
- Springel, V., Yoshida, N. & White, S. D. M. (2001), New Astronomy, 6, 79
- Stadel, J. (2000), PhD thesis, University of Washington, Seattle
- Starobinskii, A. (1982), Phys. Lett. 117B, 175
- Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. (1996), ApJ, 462, L17
- Tytler, D., O'Meara, J. M., Suzuki, N. & Lubin, D. (2000), Physica Scripta, 85, 12.
- Van Waerbeke, L. et al (2001), A&A, 374, 757.

- Viana, P. T. P. & Liddle, A. (1996), MNRAS, 281, 323
- Wilson, G., Kaiser, N., Luppino, G., & Cowie, L. L. (2001), ApJ, 555, 572
- White, S. D. M. & Frenk, C. S. (1991), ApJ, 379, 52
- White, S. D. M., Frenk, C. S. & Davis, M. (1983) ApJ 274, L1.
- White, S. D. M., Navarro, J. F., Evrard, A. E., Frenk, C. S. (1998), Nature, 366, 429.
- York, D.G, etal (the SDSS collaboration) (2000) AJ, 120, 1579
- Yoshida, N., Stoehr, F., Springel, V. & White, S. D. M. (2002), submitted to MNRAS (astro-ph/0202341)
- Zucca, E. etal (1997), AA, 326, 477.