

## THE COSMIC RADIO AND INFRARED BACKGROUNDS CONNECTION

ELI DWEK

Laboratory for Astronomy and Solar Physics, NASA Goddard Space Flight Center, Greenbelt, MD 20771;  
eli.dwek@gsfc.nasa.gov

AND

MICHAEL K. BARKER

Astronomy Department, University of Florida, Gainesville, FL 32611-2055; mbarker@astro.ufl.edu

Received 2002 January 24; accepted 2002 April 10

### ABSTRACT

We use the radio-infrared (IR) flux correlation between star-forming galaxies in the local universe to examine the connection between their cumulative contributions to the cosmic IR and radio backgrounds. The general expression relating the intensities of the two backgrounds is complicated and depends on details of the evolution of the galaxies' IR luminosity function with redshift. However, in the specific case in which the radio-IR flux correlation is linear, the relation between the intensity of the IR background and the brightness temperature of the radio background reduces to a simple analytical expression that at 178 MHz is  $I_{\text{CIB}} \text{ (nW m}^{-2} \text{ sr}^{-1}) = 2.7T_{\text{crb}} \text{ (K)}$ , where the numerical coefficient was calculated for a radio spectral index of 0.7. This relation is insensitive to the star-formation history of the galaxies that produce the cosmic IR background (CIB). We use the observed CIB intensity to constrain the cosmic star formation history and the relation between the CIB and the cosmic radio background (CRB) to constrain the relative contribution of star-forming galaxies to the CRB. Current limits on the CIB intensity predict a 178 MHz brightness temperature of  $\sim 18 \pm 9 \text{ K}$ , about half of the  $37 \pm 8 \text{ K}$  inferred for an isotropic radio component. This suggests that star-forming galaxies and active galactic nuclei (AGNs) contribute about equally to the CRB intensity at that frequency.

*Subject headings:* cosmology: observations — diffuse radiation — infrared: general —  
radio continuum: general

*On-line material:* color figures

### 1. INTRODUCTION

Star-forming galaxies exhibit a remarkable correlation between their radio and infrared (IR) fluxes, covering over four decades of IR flux intensities (Helou, Soifer, & Rowan-Robinson 1985, and references therein). The correlation results from the fact that the radio and IR fluxes are different manifestations of the physical processes associated with the life cycle of the same stellar objects (Helou & Bicay 1993; Lisenfeld, Völk, & Xu 1996). Massive stars heat the dust in the interstellar medium, form H II regions, and after their explosive deaths, accelerate particles to cosmic-ray energies, processes that, respectively, give rise to the observed galactic thermal IR, thermal radio, and synchrotron emission. The correlation may not hold for active galactic nuclei (AGNs), in which the radio emission is not associated with the life cycle of massive stars.

If the radio-IR correlation for individual star-forming galaxies in the local universe holds for all redshifts, then their cumulative contributions to the cosmic radio and IR backgrounds should be related. Haarsma & Partridge (1998, hereafter HP98) used this correlation to estimate the intensity of the cosmic radio background (CRB) that can be attributed to star-forming galaxies. At the time of their analysis, only the cosmic IR background (CIB) at wavelengths larger than  $\sim 120 \mu\text{m}$  was definitively detected by the Diffuse Infrared Background Experiment (DIRBE) and Far Infrared Spectrophotometer (FIRAS) instruments on board the *Cosmic Background Explorer (COBE)* satellite (Hauser et al. 1998; Fixsen et al. 1998). For simplicity, HP98 assumed

that all the sources giving rise to the CIB release their energy instantaneously at redshift  $z = 1$ . With this simple assumption about the formation epoch of the CIB, most of its detected intensity is determined by the  $\sim 80 \mu\text{m}$  spectral luminosity density of galaxies at the assumed redshift of its creation. Adopting a linear relation,  $S_\nu(20 \text{ cm}) \propto S_\nu(80 \mu\text{m})$ , between the radio and IR fluxes from star-forming galaxies, HP98 derived their contribution to the 40 cm CRB intensity. To compare this CRB estimate with the observed Bridle (1967) data point at 178 MHz, HP98 adopted a  $\nu^{-0.7}$  power-law spectrum for the radio sources. They derived a brightness temperature of  $T_{\text{crb}}(178 \text{ MHz}) \approx 15 \text{ K}$ , about half the value inferred from the observations. HP98 attributed the remaining CRB intensity to the contribution of AGNs, which do not contribute significantly to the CIB (Barger, Cowie, & Mushotzky 2001; Fadda et al. 2002).

In this paper we extend the analysis of HP98 to the more general case in which the cosmic star formation rate (CSFR) of the CIB sources is a general function of redshift. First, in § 2 we briefly review the various presentations of the radio-IR correlation and rederive this correlation in terms of the galaxies' 8–1000  $\mu\text{m}$  IR fluxes. Previous presentations expressed this correlation as a function of only the far-IR (FIR) fluxes, derived from the galaxies' 60 and 100  $\mu\text{m}$  detections by the *Infrared Astronomy Satellite (IRAS)*. Such a correlation may systematically underestimate the radio flux from more luminous star-forming galaxies in which an increasing fraction of the IR flux may be emitted in the shorter (12 and 25  $\mu\text{m}$ ) *IRAS* bands. We then explore the

functional form of the radio-IR correlation. HP98 adopted a linear relation between the radio and IR fluxes, which may not be an accurate fit to the data. Significant deviations from linearity will require detailed knowledge of the evolution of the galaxies' IR luminosity function with redshift. However, we argue that the data are consistent with a linear relation over most of the range of the IR luminosities that contribute to the local IR luminosity density. In § 3 we first derive the expressions for the CIB and CRB intensities produced by star-forming galaxies for a general power-law correlation between their radio and IR fluxes. We then derive a simple analytic expression relating these background intensities when their radio-IR correlation is linear. In § 4 we apply these results to different cosmic star-formation histories with their associated CIB intensity limits. A brief summary of our paper is presented in § 5.

## 2. THE RADIO-INFRARED FLUX CORRELATION

The radio-IR correlation can be written in a general form as

$$P_\nu(\nu, L_{\text{IR}})(\text{W Hz}^{-1}) = \kappa(\nu) \left( \frac{L_{\text{IR}}}{L_\odot} \right)^{\beta(\nu)}, \quad (1)$$

where  $L_{\text{IR}}$  is the IR luminosity of the galaxy,  $P_\nu(\nu)$  is the specific radio luminosity at the radio frequency  $\nu$ , and  $\kappa$  and  $\beta$  are coefficients that depend on the frequency at which the correlation is expressed. The coefficient  $\kappa$  is dimensional, and its value depends on the units used in the correlation. The presentation above is often referred to as the  $\{P_\nu-L_{\text{IR}}\}$  presentation of the radio-IR correlation. An alternative is the  $\{S_\nu-F_{\text{IR}}\}$  presentation, in which the radio and IR outputs are expressed in units of  $\text{W m}^{-2} \text{ Hz}^{-1}$  and  $\text{W m}^{-2}$ , respectively.

The two presentations are, of course, equivalent for individual galaxies. However, for determining  $\beta$ , the choice of presentation is important if  $\beta$  is a priori known to be different from unity (Cox et al. 1988). If all galaxies lie on a line of slope  $\beta = 1$  in a  $\log S_\nu$  versus  $\log F_{\text{IR}}$  plot, then the conversion to a  $\{P_\nu-L_{\text{IR}}\}$  presentation will move them up or down along a diagonal line, leaving the overall slope of the correlation unchanged. However, if the slope of the correlation is different from unity in a  $\{S_\nu-F_{\text{IR}}\}$  presentation, then fainter galaxies, which tend to be more distant than brighter ones, may move systematically away or toward the  $\beta = 1$  slope line in a  $\{P_\nu-L_{\text{IR}}\}$  presentation. The overall effect will be to flatten the slope if  $\beta < 1$  and to steepen it if  $\beta > 1$ , in a transition from a  $\{S_\nu-F_{\text{IR}}\}$  to a  $\{P_\nu-L_{\text{IR}}\}$  presentation.

Various investigators examined the radio-IR correlation using a variety of sample selection criteria. All used FIR fluxes measured by the *IRAS* satellite and constructed galaxy FIR luminosities (fluxes) from the 60 and 100  $\mu\text{m}$  detections using the following relation (Sanders & Mirabel 1996),

$$\begin{aligned} F_{\text{FIR}}(\text{W m}^{-2}) &= 1.26 \times 10^{-14} \\ &\times [2.58S_\nu(60 \mu\text{m}) + S_\nu(100 \mu\text{m})], \\ L_{\text{FIR}}(L_\odot) &= 1.26 \times 10^{12} [2.58L_\nu(60 \mu\text{m}) + L_\nu(100 \mu\text{m})], \end{aligned} \quad (2)$$

where  $S_\nu$  and  $L_\nu$  are flux and luminosity densities expressed in units of  $\text{Jy}$  ( $=10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$ ) and  $L_\odot \text{ Hz}^{-1}$ , respectively.

As we show in § 3, the value of  $\beta$  plays an important role in quantifying the CIB-CRB connection. A value of  $\beta$  that is significantly different from unity will require detailed knowledge of the evolution of the IR luminosity function with redshift, whereas a value of unity will considerably simplify the relation between the two background emissions. We therefore briefly review previous determinations of  $\beta$ .

Initial investigations, using a relatively small sample of 91 radio-selected galaxies observed at  $\nu = 4.8 \text{ GHz}$  (de Jong et al. 1985) and 38 optically and radio-selected galaxies observed at 1.4 GHz (Helou et al. 1985), found the radio-IR correlation to be linear. Wunderlich & Klein (1988) extended the analysis of de Jong et al. (1985) to a wider range of galaxy types, covering about 4 orders of magnitude in FIR luminosities. They found that the correlation at  $\nu = 4.8 \text{ GHz}$  is approximately linear ( $\beta = 0.99 \pm 0.38$ ) up to IR luminosities of  $9.1 \times 10^9 L_\odot$ , with  $\beta$  increasing to a value of  $1.26 \pm 0.33$  at higher luminosities. However, we found that their sample of galaxies could also be fitted with a single power law with a slope of  $\beta = 1.05 \pm 0.03$ .

Devereux & Eales (1989) correlated the 1.49 GHz power with the FIR output from an optically selected sample of galaxies with luminosities between  $\approx 10^8$  and  $10^{11} L_\odot$ . They found the slope of the correlation to be significantly larger than unity with a value of  $\beta = 1.28$ . A similar conclusion was reached by Cox et al. (1988), who studied the correlation for a flux-limited sample of 74 radio galaxies at 151 MHz. They found  $\beta = 1.32 \pm 0.06$  in a  $\{S_\nu-F_{\text{IR}}\}$  presentation and  $\beta = 1.15 \pm 0.04$  in a  $\{P_\nu-L_{\text{IR}}\}$  presentation of the data and adopted an average slope of  $\beta = 1.23$ .

Chi & Wolfendale (1990) provided theoretical arguments for why the slope of the radio-IR correlation should not be unity. They predicted a break in the slope of the correlation at an IR luminosity above which most of the electrons producing the radio synchrotron emission remained trapped in the galaxy. For these galaxies, the slope of the correlation should be unity, but for the smaller and less IR luminous galaxies, they argued that  $\beta$  should be larger than unity. They claimed to have found evidence for this effect in the data and reported values of  $\beta = 1.37$  for low-luminosity galaxies, a trend exactly opposite to that found by Wunderlich & Klein (1988). Our analysis of the Chi & Wolfendale (1990) sample did not confirm either trend. We found that their data could be represented by a single slope with a value of  $\beta = 1.27 \pm 0.06$ .

Condon, Anderson, & Helou (1991, hereafter CAH91) reexamined the radio-IR correlation using galaxies from the *IRAS* revised bright galaxy sample (BGS) that were detected with the Very Large Array at 1.49 GHz and subtracted from the sample spectroscopically identified AGNs. They found the slope of the correlation to be significantly greater than unity with a smaller value of  $\beta = 1.11 \pm 0.02$  over the  $\sim 10^9$ – $10^{13} L_\odot$  luminosity range.

For the purpose of the present analysis, we reexamined the radio-IR correlation starting from the same galaxy sample used by CAH91, the revised *IRAS* BGS, eliminating galaxies that were not detected in all four *IRAS* bands and removing AGNs using a more recent catalog of spectroscopically confirmed AGN galaxies (Véron-Cetty & Véron 2001).<sup>1</sup> The total number of galaxies left in the sample is

<sup>1</sup> See A Catalogue of Quasars and Active Nuclei (10th ed.; St. Michel l'Observatoire: Haute-Provence Obs.).




FIG. 1.—Radio fluxes of galaxies in the *IRAS* BGS, plotted vs. their 8–1000  $\mu\text{m}$  luminosity determined from eq. (3). Star-forming galaxies are represented by crosses, and spectroscopically confirmed AGNs by open triangles. The lines are fits to the radio-IR correlation between the star-forming galaxies only. The thin line represents the best power-law fit, and the thick line the best linear fit to the data. The population of AGNs generally follows the same correlation, but with a wider dispersion. [See the electronic edition of the *Journal* for a color version of this figure.]

222, compared to 258 galaxies used in the analysis of CAH91. We used the additional 12 and 25  $\mu\text{m}$  *IRAS* bands to derive the total 8–1000  $\mu\text{m}$  IR luminosity from the relation (Sanders & Mirabel 1996)

$$L_{\text{IR}}(L_{\odot}) = 1.8 \times 10^{12} [13.48L_{\nu}(12 \mu\text{m}) + 5.16L_{\nu}(25 \mu\text{m}) + 2.58L_{\nu}(60 \mu\text{m}) + L_{\nu}(100 \mu\text{m})]. \quad (3)$$

Figure 1 depicts the radio-IR correlation for a  $\{P_{\nu}-L_{\text{IR}}\}$  presentation of the data and the best power-law fit to the data. We derive an almost identical slope  $\beta$  for the same sample of galaxies as CAH91, with a smaller value for  $\kappa$ , reflecting the fact that the 8–40  $\mu\text{m}$  emission was not included in their galactic IR energy budget. The values of the fit for radio-IR correlation are  $\{\kappa, \beta\} = \{2.96 \times 10^{10}, 1.083 \pm 0.02\}$ . An almost identical slope was obtained for the  $\{S_{\nu}-F_{\text{IR}}\}$  presentation, suggesting that there were no significant flux-related systematic errors in determining the distances of the galaxies in the sample. Also shown in Figure 1 is a forced linear fit to the data with  $\{\kappa, \beta\} = \{2.47 \times 10^{11}, 1.0\}$ . The linear fit is almost indistinguishable from the best fit for galaxies with luminosities above  $\sim 10^{10} L_{\odot}$ , which produce most of the IR luminosity density in the local and high- $z$  universe. The same linear fit would provide a similarly good representation of the radio-IR correlation even if the AGNs were included in the data. The AGN population, however, shows more dispersion around the fit.

### 3. THE CIB AND CRB INTENSITIES PRODUCED BY STAR-FORMING GALAXIES

The specific intensity  $I_{\nu}^{\text{crb}}(\nu_R)$  of the CRB at the observed radio frequency  $\nu_R$  is given by the integral over sources (e.g., Peebles 1993)

$$I_{\nu}^{\text{crb}}(\nu_R) = \left(\frac{c}{4\pi}\right) \int_0^{\infty} \mathcal{E}_{\nu}(\nu, z) \left| \frac{dt}{dz} \right| dz, \quad (4)$$

where  $\mathcal{E}_{\nu}(\nu, z)$  is the specific radio luminosity per comoving volume element at redshift  $z$ ,  $\nu = \nu_R(1+z)$  is the frequency in the rest frame of the luminous objects, and  $|dt/dz|$  is given by (Hogg 1999)

$$\left| \frac{dt}{dz} \right|^{-1} = H_0(1+z) \left[ (1+z)^2(1+\Omega_m z) - z(2+z)\Omega_{\Lambda} \right]^{1/2}, \quad (5)$$

where  $H_0$  is the Hubble constant,  $h$  is its value in units of  $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_m \equiv \rho_m/\rho_c$  is the present mass density of the universe normalized to the critical density  $\rho_c = 1.88 \times 10^{-29} h^2 \text{ g cm}^{-3}$ , and  $\Omega_{\Lambda} \equiv \Lambda/3H_0^2$  is the dimensionless cosmological constant.

We assume that the radio-IR correlation, expressed at a given rest-frame frequency  $\nu_0$ , holds for all redshifts and that the radio spectrum of the individual galaxies follows a  $\sim \nu^{-\alpha}$  power law with spectral index  $\alpha$ , so that  $P_{\nu}(\nu, L_{\text{IR}}, z) = (\nu/\nu_0)^{-\alpha} P_{\nu}(\nu_0, L_{\text{IR}}, z)$ . The spectral luminosity density  $\mathcal{E}_{\nu}$  can then be expressed as

$$\begin{aligned} \mathcal{E}_{\nu}(\nu, z) &= \int_{L_{\text{min}}}^{L_{\text{max}}} P_{\nu}(\nu, L_{\text{IR}}, z) \Phi(L_{\text{IR}}, z) dL_{\text{IR}} \\ &= \kappa_0 \left(\frac{\nu}{\nu_0}\right)^{-\alpha} \int_{L_{\text{min}}}^{L_{\text{max}}} L_{\text{IR}}^{\beta_0} \Phi(L_{\text{IR}}, z) dL_{\text{IR}}, \end{aligned} \quad (6)$$

where  $\Phi(L_{\text{IR}}, z) \equiv [d\varphi(L_{\text{IR}}, z)/dL_{\text{IR}}]$  is the differential IR luminosity function,  $d\varphi(L_{\text{IR}}, z)$  is the comoving number density of galaxies with luminosities between  $L_{\text{IR}}$  and  $L_{\text{IR}} + dL_{\text{IR}}$  in the  $\{L_{\text{min}}, L_{\text{max}}\}$  luminosity interval, and  $\kappa_0$  and  $\beta_0$  are the parameters of the radio-IR correlation at frequency  $\nu_0$ .

The specific intensity of the CRB can now be written as

$$I_{\nu}^{\text{crb}}(\nu_R) = \left(\frac{c}{4\pi}\right) \left(\frac{\nu_R}{\nu_0}\right)^{-\alpha} \int_0^{\infty} \frac{f(z) \mathcal{L}_{\text{IR}}(z)}{(1+z)^{\alpha}} \left| \frac{dt}{dz} \right| dz, \quad (7)$$

where  $\mathcal{L}_{\text{IR}}(z)$  is the comoving IR luminosity density at redshift  $z$ ,

$$\mathcal{L}_{\text{IR}}(z) \equiv \int_{L_{\text{min}}}^{L_{\text{max}}} L_{\text{IR}} \phi(L_{\text{IR}}, z) dL_{\text{IR}}, \quad (8)$$

and  $f(z)$  is defined as

$$f(z) \equiv \frac{\kappa_0 \int_{L_{\text{min}}}^{L_{\text{max}}} L_{\text{IR}}^{\beta_0} \phi(L_{\text{IR}}, z) dL_{\text{IR}}}{\mathcal{L}_{\text{IR}}(z)}. \quad (9)$$

The same star-forming galaxies give rise to the CIB with an intensity  $I_{\text{CIB}}$  that is given by

$$I_{\text{CIB}} = \left(\frac{c}{4\pi}\right) \int_0^{\infty} \frac{\mathcal{L}_{\text{IR}}(z)}{1+z} \left| \frac{dt}{dz} \right| dz. \quad (10)$$

For a linear correlation,  $\beta_0 = 1$  and  $f(z) = \kappa_0 (\text{Hz}^{-1})$ . The spectral intensity of the CRB then becomes

$$I_{\nu}^{\text{crb}}(\nu_R) = \left(\frac{c}{4\pi}\right) \left(\frac{\nu_R}{\nu_0}\right)^{-\alpha} \kappa_0 \int_0^{\infty} \frac{\mathcal{L}_{\text{IR}}(z)}{(1+z)^{\alpha}} \left| \frac{dt}{dz} \right| dz. \quad (11)$$

In this case, the relation between  $I_{\nu}^{\text{crb}}(\nu_R)$  and  $I_{\text{CIB}}$  is consid-

erably simplified and is given by

$$\begin{aligned} I_{\text{CIB}} &= \left(\frac{\nu_R}{\nu_0}\right)^\alpha \frac{g(\alpha)}{\kappa_0} I_{\nu}^{\text{crb}}(\nu_R) \\ &= 1.18 \times 10^8 \left(\frac{\nu_R}{\nu_0}\right)^\alpha \frac{g(\alpha)}{\kappa_0} \\ &\quad \times \nu_R^2 (\text{MHz}) T_{\text{crb}}(\nu_R) \text{ nW m}^{-2} \text{ sr}^{-1}, \end{aligned} \quad (12)$$

where

$$\begin{aligned} g(\alpha) &\equiv \left[ \int_0^\infty \frac{\mathcal{L}_{\text{IR}}(z)}{1+z} \left| \frac{dt}{dz} \right| dz \right] \\ &\quad \times \left[ \int_0^\infty \frac{\mathcal{L}_{\text{IR}}(z)}{(1+z)^\alpha} \left| \frac{dt}{dz} \right| dz \right]^{-1}, \end{aligned} \quad (13)$$

and  $T_{\text{crb}}(\nu_R)$  is the CRB brightness temperature at frequency  $\nu_R$ .

#### 4. THE CIB-CRB CONNECTION FOR VARIOUS COSMIC STAR FORMATION HISTORIES

Several models have been recently developed for calculating the evolution of  $\mathcal{L}_{\text{IR}}(z)$  with redshift. Here we focus on some select representative cases: Malkan & Stecker (2001; baseline model), Chary & Elbaz (2001; pure luminosity evolution model), and Xu et al. (2001; peak model). These models were developed to explain the limits and detections of the CIB spectrum and galaxy number counts obtained with the *IRAS*, the *Infrared Space Observatory*, and the Submillimeter Common-User Bolometric Array instrument on the James Clerk Maxwell Telescope at various IR and submillimeter wavelengths. Figure 2 presents the evolution of  $\mathcal{L}_{\text{IR}}(z)$  for these select models. Also plotted are observational estimates of the comoving CSFR at different redshifts. Star formation rates were converted to IR luminosity densities using the relation (Kennicutt 1998)




FIG. 3.—Function  $g(\alpha)$ , defined by eq. (13), plotted against  $\alpha$  for the various CSFRs depicted in Fig. 2. The thick line represents a  $\alpha^{0.688}$  power-law fit to the function for  $\alpha$  values between 0.6 and 1.2. [See the electronic edition of the *Journal* for a color version of this figure.]

$\mathcal{L}_{\text{IR}} (L_\odot \text{ Mpc}^{-3}) = 6 \times 10^9 \rho_*$  ( $M_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ ). References to the CSFR data can be found in the papers listed earlier. Also shown in the figure are approximate upper and lower limits to the IR luminosity density. The top-hat function centered on redshift  $z = 1$  in the figure represents the instantaneous energy injection model used by HP98 to calculate the CRB. The amplitude of the function was chosen so that it reproduced the nominal  $\sim 5\text{--}1000 \mu\text{m}$  CIB intensity of  $50 \text{ nW m}^{-2} \text{ sr}^{-1}$  (Hauser & Dwek 2001).

Figure 3 shows the value of  $g(\alpha)$  as a function of  $\alpha$  for the different star formation histories depicted in Figure 2. The function  $g(\alpha)$  is insensitive to the cosmic star formation history, a direct consequence of the fact that the integrands in equation (13) differ by only a factor of  $(1+z)^{\alpha-1}$ . For  $\alpha \approx 0.6\text{--}1.2$ , the most probable range of values for the radio spectral index,  $g(\alpha)$  is well approximated by

$$g(\alpha) = \alpha^{0.688} \quad \text{for } \alpha \approx 0.6\text{--}1.2 \quad (14)$$

to an accuracy of better than 2%.

Using equations (12) and (14) and the linear radio-IR correlation at  $\nu_0 = 1.49 \text{ GHz}$ ,  $P_\nu (\text{W Hz}^{-1}) = 2.47 \times 10^{11} L_{\text{IR}} (L_\odot)$ , we can express the CIB intensity in terms of the radio brightness temperature at frequency  $\nu_R = 178 \text{ MHz}$  as

$$\begin{aligned} I_{\text{CIB}} (\text{nW m}^{-2} \text{ sr}^{-1}) &= \mathcal{A}(\alpha) T_{\text{crb}}, \\ \text{where } \mathcal{A}(\alpha) &= \{3.0, 2.7, 2.4, 2.1\} \\ \text{for } \alpha &= \{0.6, 0.7, 0.8, 0.9\}. \end{aligned} \quad (15)$$

Table 1 lists the values of the CIB intensity and the 178 MHz brightness temperature obtained from the respective use of equations (10) and (11) for the different star formation histories and radio spectral index  $\alpha$ . The entries in the table satisfy the analytical approximation for the relation between  $I_{\text{CIB}}$ ,  $T_{\text{crb}}$ , and  $\alpha$  to an accuracy of a few percent. For comparison, we also list the observed limits and detections of the CIB (Hauser & Dwek 2001) and the 178 MHz CRB brightness temperature inferred from the Bridle data (Bridle 1967) for the different values of  $\alpha$ . Also shown in the table are the observed  $T_{\text{crb}}/I_{\text{CIB}}$  ratios and those predicted by the CIB-CRB correlation (eq. [15]).

It is interesting to compare our model predictions with the result obtained by HP98. HP98 found that the star-




FIG. 2.—Comoving IR luminosity density predicted by several galaxy number count models as a function of redshift. The data points represent optical and IR determinations of the comoving CSFR as a function of redshift (references to the data can be found in the papers listed in § 4). A scaling factor of  $6 \times 10^9$  was used to convert the CSFR (in  $M_\odot \text{ yr}^{-1}$ ) to an IR luminosity density (in  $L_\odot \text{ Mpc}^{-3}$ ). The top and bottom dashed lines represent approximate upper and lower limits to the CSFR. [See the electronic edition of the *Journal* for a color version of this figure.]

TABLE 1  
CIB AND CRB INTENSITIES PREDICTED BY VARIOUS MODELS FOR THE COSMIC STAR FORMATION RATE<sup>a</sup>

MODEL	$I_{\text{CIB}} (3.5\text{--}1000 \mu\text{m})$ ( $\text{nW m}^{-2}\text{sr}^{-1}$ )	$T_{\text{crb}}$ AT 178 MHz (K)			
		$\alpha = 0.6$	$\alpha = 0.7$	$\alpha = 0.8$	$\alpha = 0.9$
Minimum CSFR.....	8.5	2.7	3.1	3.5	4.1
Maximum CSFR.....	73	24.8	28.0	31.5	35.6
Haarsma & Partridge 1998.....	50	15.7	18.1	20.8	24.1
Malkan & Stecker 2001.....	23	8.2	9.1	10.2	11.4
Chary & Elbaz 2001.....	37	12.3	13.9	15.8	18.0
Xu et al. 2001.....	58	18.8	21.5	24.5	27.9
Observational limits <sup>b</sup> .....	$50 \pm 25$	$57 \pm 11$	$37 \pm 8$	$23 \pm 5$	$15 \pm 3$
$\mathcal{A}^{-1} \equiv T_{\text{crb}}/I_{\text{CIB}}$ (calculated) <sup>c</sup> .....	...	0.33	0.37	0.42	0.48
$T_{\text{crb}}/I_{\text{CIB}}$ (observed).....	...	$1.1 \pm 0.6$	$0.7 \pm 0.4$	$0.5 \pm 0.25$	$0.30 \pm 0.16$

<sup>a</sup> The CSFRs predicted by the tabulated models are shown in Fig. 2. CIB intensities and CRB temperatures were calculated from eqs. (10) and (11), respectively. The parameter  $\alpha$  is the spectral index of the radio sources.

<sup>b</sup> Observational limits on the CIB intensity are summarized in Hauser & Dwek 2001. CRB temperatures were taken from Bridle 1967: Table VIII.

<sup>c</sup> Calculated using eq. (15).

forming galaxies that produce the detected 120–260  $\mu\text{m}$  CIB intensity of  $\sim 22 \text{ nW m}^{-2} \text{ sr}^{-1}$  contribute about 15 K to the 178 MHz brightness temperature (they adopted a value of  $\alpha = 0.7$ ). This result seems at first glance to be in disagreement with the entry for the HP98 model in the table. The reason for this apparent “discrepancy” is that HP98 expressed the radio-IR correlation in terms of the FIR luminosity of galaxies. Had they expressed this correlation in terms of their IR luminosity, which is about twice the FIR value, they would have derived a radio brightness temperature of  $\sim 17$  K for a 3.5–1000  $\mu\text{m}$  background intensity of  $50 \text{ nW m}^{-2} \text{ sr}^{-1}$ , almost identical to the value listed in the table. The analytical expression presented in this paper reproduces the results derived by HP98 for their specific CIB production scenario and generalizes their treatment to any cosmic star formation history.

## 5. DISCUSSION AND SUMMARY

In this paper we derived an analytical expression for the correlation between the CIB and CRB intensities as a function of  $\alpha$ , the spectral index of the radio sources (see eq. [15]). This correlation is summarized for select galaxy number counts and CIB models in Table 1. Several conclusions can be drawn from a simple examination of the table: (1) The minimal CSFR, defined by the lower envelope of the data, is definitely ruled out as a viable representation of the CSFR, since it falls short of providing the observed intensity of the CIB. The maximal CSFR, defined by the upper envelope of the data, is consistent with the CIB and CRB limits

only for values of  $\alpha \lesssim 0.8$ . (2) The baseline model of Malkan & Stecker (2001) is barely consistent with the lower limit on the CIB intensity, and consequently, for  $\alpha = 0.7$ , it requires AGNs to contribute more than 70% of the 178 MHz background. (3) For  $\alpha = 0.9$ , all models, with the exception of those of Malkan & Stecker (2001) and Chary & Elbaz (2001), predict CRB temperatures that are  $2 \sigma$  above the observed value. The latter two models leave no room for a significant contribution of AGNs to the CRB. (4) For values of  $\alpha \approx 0.6\text{--}0.8$ , the calculated  $T_{\text{crb}}/I_{\text{CIB}}$  ratio is lower than the nominal observed ratios, suggesting that respectively  $\sim 70\%$  to  $\sim 16\%$  of the CRB at 178 MHz must arise from AGNs, regardless of the radiative history of star-forming galaxies.

All conclusions listed above assume that the radio-IR correlation observed in the local universe can be extended to galaxies at all redshifts. Future observations will provide a larger sample of galaxies with which to study the radio-IR correlation and place tighter limits on the CIB, advances that will lead to a better understanding of the relative contribution of star-forming galaxies and AGNs to the CRB.

We thank Rick Arendt for useful discussions and his insightful comments on an earlier version of the manuscript. E. D. acknowledges NASA’s Astrophysics Theory Program NRA 99-OSS-01 for support of this work. M. B.’s summer student internship at NASA/GSFC was supported by the “Research Opportunities for Undergraduates in the Laboratory for Astronomy and Solar Physics” program.

## REFERENCES

- Barger, A. J., Cowie, L. L., & Mushotzky, R. F. 2001, *AJ*, 121, 662  
 Bridle, A. H. 1967, *MNRAS*, 136, 219  
 Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562  
 Chi, X., & Wolfendale, A. W. 1990, *MNRAS*, 245, 101  
 Condon, J. J., Anderson, M. L., & Helou, G., 1991, *ApJ*, 376, 95 (CAH91)  
 Cox, M. J., Eales, S. A. E., Alexander, P., & Fitt, A. J. 1988, *MNRAS*, 235, 1227  
 de Jong, T., Klein, U., Wielebinski, R., & Wunderlich, E. 1985, *A&A*, 147, L6  
 Devereux, N. A., & Eales, S. A. 1989, *ApJ*, 340, 708  
 Fadda, D., Flores, H., Hasinger, G., Franceschini, A., Altieri, B., Cesarsky, C. J., Elbaz, D., & Ferrando, Ph. 2002, *A&A*, 383, 838  
 Fixsen, D. J., Dwek, E., Mather, J. C., Bennett, C. L., & Shafer, R. A. 1998, *ApJ*, 508, 123  
 Haarsma, D. B., & Partridge, R. B. 1998, *ApJ*, 503, L5 (HP98)  
 Hauser, M. G., & Dwek, E. 2001, *ARA&A*, 39, 249  
 Hauser, M. G., et al. 1998, *ApJ*, 508, 25  
 Helou, G., & Bica, M. D. 1993, *ApJ*, 415, 93  
 Helou, G., Soifer, B. T., & Rowan-Robinson, M. 1985, *ApJ*, 298, L7  
 Hogg, D. W. 1999, preprint (astro-ph/9905116)  
 Kennicutt, R. C., Jr. 1998, *ARA&A*, 36, 189  
 Lisenfeld, U., Völk, H. J., & Xu, C. 1996, *A&A*, 306, 677  
 Malkan, M. A., & Stecker, F. W. 2001, *ApJ*, 555, 641  
 Peebles, P. J. E. 1993, *Principles of Physical Cosmology* (Princeton: Princeton Univ. Press)  
 Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749  
 Véron-Cetty, M. P., & Véron, P. 2001, *A&A*, 374, 92  
 Wunderlich, E., & Klein, U. 1988, *A&A*, 206, 47  
 Xu, C., Lonsdale, C. J., Shupe, D. L., O’Linger, J., & Masci F. 2001, *ApJ*, 562, 179