CSIRO PUBLISHING

www.publish.csiro.au/journals/pasa *Publications of the Astronomical Society of Australia* 0000, 000, 000–000

Large Scale Structure in the Local Universe – The 2MASS Galaxy Catalog

Thomas Jarrett,[1,2]

[1]Infrared Processing and Analysis Center, MS 100-22, California Institute of Technology, Pasadena, CA 91125
[2]jarrett@ipac.caltech.edu

Abstract
Using twin ground-based telescopes, the Two Micron All Sky Survey (2MASS) scanned both equatorial hemispheres, detecting more than 500 million stars and resolving more than 1.5 million galaxies in the near-infrared (1 - 2.2 μm) bands. The Extended Source "Galaxy" Catalog (XSC) embodies both photometric and astrometric whole sky uniformity, revealing large scale structures in the local Universe and extending our view into the Milky Way's dust-obscured "Zone of Avoidance." The XSC represents a uniquely unbiased sample of "normal" galaxies, particularly sensitive to the underlying, dominant, stellar mass component of galaxies. The basic properties of the XSC, including photometric sensitivity, source counts, and spatial distribution are presented here. **We illustrate the spatial clustering properties, ranging from interacting galaxies to groups and clusters, and finally to the largest "cosmic web" of superclusters spanning the sky.**

Keywords: galaxies: statistics-galaxies: photometry-galaxies: general-galaxies: fundamental parameters-infrared: galaxies-clusters: surveys-astronomical databases: atlases

30 March 2004



Figure 1. Panoramic view of the entire near-infrared sky reveals the distribution of galaxies beyond the Milky Way. The image is derived from the 2MASS Extended Source Catalog (XSC) – more than 1.5 million galaxies, and the Point Source Catalog (PSC) – nearly 0.5 billion Milky Way stars. The galaxies are color coded by "redshift," photometrically deduced from the K band (2.2 μm) or as given in the NASA Extragalactic Database (NED). Blue are the nearest sources ($z < 0.01$); green are at moderate distances ($0.01 < z < 0.04$) and red are the most distant sources that 2MASS resolves ($0.04 < z < 0.1$). The map is projected with an equal area aitoff in the Galactic system (Milky Way at center). A key to this projection is given in Figure 2.

© Astronomical Society of Australia 0000



Figure 2. Galactic projection of the 2MASS Galaxy Catalog. Galaxy clusters and large scale structures are labeled. The complete projection in shown in Figure 1.



Figure 3. Total number of galaxies per deg^2 per mag internal for $|b| > 25°$ (total area: ∼24,000 deg^2). The J, H and Ks source counts are represented with blue, green and red lines, respectively. For comparison, the narrow (but deep) galaxy counts of Glazebrook et al (1994) and Gardner et al (1997) are shown in black.

1 Introduction

Our understanding of the origin and evolution of the Universe has been fundamentally transformed with seminal redshift, distant supernovae and cosmic microwave background surveys. The focus has shifted to the distribution and nature of dark matter and dark energy that drive the dynamics of the expanding cosmos. The study of the local Universe, including its peculiar motions and its clustering on scales exceeding 100 Mpc, is an essential ingredient in the connection between the origin of structure in the early Universe and the subsequent formation of galaxies and their evolution to the state we observe today. Key issues include the location and velocity distribution of galaxies, leading to the mass-to-light relationship between what is observed and what is influencing the mass density field.

Spurred on by the enormous success of the redshifts surveys (cf. Huchra et al 1982), conducted some 20 years ago, pioneers such as CfA's John Huchra proposed to image the entire sky at near-infrared wavelengths to create an unbiased census of galaxies that would fuel the next generation of redshift surveys (e.g., 6dFGS; see Watson et al 2001 and Jones et al 2004) and help address these *weighty* issues. After much deliberation, what finally transpired some ten years ago now, is the Two Micron All Sky Survey (2MASS). Many years of observations and hard work to detect and extract sources has produced a gap-free image atlas of entire sky, and catalogues containing stars and galaxies. The 2MASS view of the sky–the spatial distribution of galaxies in the local Universe–is the focus of this paper. Figure 1 is an attempt to encapsulate our present understanding of the large scale structure that embodies the local Universe.

2 Two Micron All Sky Survey

2MASS is a ground-based, near-infrared survey of the whole sky. It began operations in the mid-1990's and completed observations in early 2001. The final source catalogs and image Atlas was released to the public in the fall of 2002 (Cutri et al 2000). 2MASS is fundamentally an imaging survey, with detection and source characterization derived from images that span the near-infrared windows: J (1.2 μm), H (1.6 μm), and Ks (2.2 μm). The images were acquired using an efficient drift scan and freeze-frame technique, painting the sky with $8.5' \times 6°$ tiles or "scans" (Skrutskie et al 1997), forming 23 separate images per tile per near-infrared band of size 512 × 1024 pixels with resampled $1''$ pixels. A total of 4,121,439 FITS images cover 4πsteradians of the sky. These images, $8.5' \times 17'$ in angular size, are also known as "coadds" since they are comprised of ∼6 optimally dithered samples per pixel. The effective beam or PSF FWHM is ∼2 to $3''$, depending on the atmospheric seeing, and is roughly the same for each band. The typical 1-sigma background noise is 21.4, 20.6, and 20.0 mag $arcsec^{-2}$ for J, H and Ks, respectively. The images include a photometric zero-point calibration that is accurate to $2-3\%$ and an astrometric solution that is accurate to $<0.2''$ (Cutri et al 2000).

Stars and galaxies are detected and characterized from the 2MASS images. The Point Source Catalog (PSC) contains ∼500 million objects, largely comprised of stars from the Milky Way. The Extended Source Catalog (XSC) contains ∼1.6 million objects clearly resolved by 2MASS, chiefly comprised of extragalactic sources in the local Universe. This paper will focus on these resolved galaxies. The images and source catalogs are available to the public *via* the Infrared Science Archive (IRSA) and NASA Extragalactic Database (NED) of IPAC.

© Astronomical Society of Australia 0000



Figure 4. Cumulative number of galaxies per deg^2 with Galactic latitude. Three Ks-band flux limits are shown: 12.5 (black), 13.0 mag (green) and 13.5 mag (blue). All longitudes are used to draw the galaxy sample. For comparison, the integrated counts per deg^2 for a field deep in the ZoA (l=50°, b= 2°) is indicated with a green triangle.

3 The 2MASS Galaxy Catalog

The extended source catalog (XSC) contains over 1.6 million spatially resolved astronomical sources: primarily (> 98%) galaxies, and to a lesser extent Galactic diffuse nebulae, H II regions, stellar clusters, planetary nebulae and young stellar objects. The *galaxy catalog* was constructed to satisfy the survey science requirements (Jarrett et al 2000), most important being the reliability (∼ 99%) and completeness (> 90%) for unconfused regions of the sky, $|b| > 20°$. These requirements are achieved for sources brighter than Ks = 13.5 mag (∼2.7 mJy) and resolved diameters larger than ∼10-15 arcsec. The differential source counts, Figure 3, illustrate the depth and areal coverage of the 2MASS galaxy catalog for unobscured regions of the sky. The Ks-band counts are steady up to ∼14 mag, with the faint end supported by the more sensitive J-band observations (complete down to 15.3 mag).

Morphology – As a function of Hubble or morphological type, 2MASS is most sensitive to early-type spirals and ellipticals (whose light is dominated by the older population of stars emitting in the near-infrared), and less sensitive to late-type spirals (whose light is dominated by the younger, hotter disk population), dwarfs (low surface brightness) and compact objects (resolution limitations of 2MASS); see Jarrett (2000) and Jarrett et al (2003). Consequently, the 2MASS galaxy catalog is partial to *normal* "old" galaxies, which are typically lower in mass-to-light compared to gas-rich spiral galaxies (see section 4 below).

Zone of Avoidance – A primary science driver of 2MASS was to penetrate the dust mask of the Milky Way. At near-infrared wavelengths, the opacity of dust is significantly smaller (∼1/10) compared to that of optical wavelengths, and thus amenable to penetrating observations. The Galactic "zone of avoidance" (ZoA) is still, however, a formidable barrier due to the sheer number of stars that block background light as well as produce a foreground (confusion) "noise." Near the center of the Milky Way the confusion noise is extreme, blocking nearly 100% of the background light; whereas far from the Galactic center the confusion noise is minimal and the veil of the Milky Way is lifted at near-infrared wavelengths. As is shown in Figure 4, 2MASS penetrates deep into the ZoA, extracting galaxies well down to $|b|$ ∼5 degrees, where detection completeness declines by ∼0.5 mag (Jarrett et al 2000).

All Sky Galaxy Distribution – Simply by counting the number of galaxies along the line of sight it is a straightforward exercise to create a crude map of the local Universe. Spatial over-densities from galaxy clusters trace the large scale structure–see for example the beautiful maps of Courtois et al (2004) who constructed the extragalactic sky using galaxies archived in LEDA. A more constructive way to map the surface density of galaxies is to integrate the 2-micron flux from all galaxies along the line of sight, thereby weighting the nearest structures and producing contrast between the Local Supercluster (cf. Tully 1982; Tonry 2000) and the more distant "cosmic web" structures; see Figure 5. This technique mitigates the biasing effects of non-uniform incompleteness due to surface brightness differences and galaxy morphology (see above). The figure illustrates how 2MASS creates a uniform view of the local Universe, except for the extreme Galactic Center, bridging the two hemispheres above and below the plane of the Milky Way (center region of figure).

A further enhancement to the all sky maps is to color-code the galaxies according to their total integrated flux. Since the integrated flux is strongly correlated with the distance to the object (assuming 2MASS galaxies have roughly the same luminosity; see the next section), the color-coding effectively adds depth to the surface density maps. This is a "poor man's" redshift, which conveniently provides a qualitative view of the 3-D galaxy distribution, illustrated in Figure 6 using a Supergalactic projection. This crude but effective method delineates real large scale structure in the local Supercluster and beyond (Figure 6). But we can go one step further by combining the integrated flux with the measured galaxy near-infrared colors to derive a photometric redshift, described in the next section.

4 The Local Universe

The current generation of large and uniform redshift surveys (e.g., CfA-RS, 2dF, SDSS, 6dFGS) provide a means to construct the 3-D space density of galaxies for volumes vastly exceeding those of the pioneering works from the 1980's. In the coming years new "radial velocity machines," such as *Echnida* (Moore, Gillingham & Saunders 2002), have the potential to increase redshifts catalogs by an order of magnitude. Nonetheless, imaging surveys, including 2MASS, SWIRE, GOODS, and in the coming years, WISE (Eisenhard & Wright 2003) catalog far more sources than the redshift surveys can ever handle. Velocity measurements will always be in the position of catching up with broad-band imaging. The interim solution to this phase-lag is to derive a "photometric redshift" using broad-band

© Astronomical Society of Australia 0000



Figure 5. Equatorial view of the 2MASS galaxy catalog (6hr RA at center). The grey-scale represents the total integrated flux along the line of sight – the nearest (and therefore brightest) galaxies produce a vivid contrast between the Local Supercluster (center-left) and the more distant cosmic web. The dark band of the Milky Way clearly demonstrates where the galaxy catalog becomes incomplete due to source confusion. Some well known large-scale structures are indicated: P-P = Perseus-Pisces supercluster, H-R = Horologium-Reticulum supercluster, P-I = Pavo-Indus supercluster, GA = "Great Attractor", GC = Galactic Center, S-C = Shapley Concentration, O-C = Ophiuchus Cluster, Virgo/Coma/Hercules = Virgo-Coma-Hercules superclusters. A 3-dimensional, Galactic projection, version of the 2MASS galaxy catalog is shown in Figure 1.

photometry measurements. Depending on the number (and spectral location) of bands used to construct the spectral energy distribution of the target galaxy, the redshift accuracy of this method ranges from 10-20% for most normal galaxies.

Photometric Redshifts – The 3-band near-infrared photometry of 2MASS is used to estimate distances to galaxies. Although this technique is very crude in terms of accuracy, it does provide a means to generate qualitative maps of the spatial distribution of galaxies and thereby construct an all sky "big picture" view of the local Universe. Here we adopt the technique devised by Jarrett et al (1998) and Kochanek et al (2003). The fundamental assumptions of this method are that galaxies have roughly (1) the same luminosity and (2) their near-infrared colors are modified by cosmic reddening (Figure 7). This method is particularly adept at revealing galaxy clusters since the redshift uncertainty declines with the sqrt of the number of cluster members detected by 2MASS. The photometric redshift of galaxy clusters is typically accurate to $\sim$ 20% using only 2MASS photometry.

By assuming that galaxies are standard candles, the distance or redshift is derived from the integrated flux, distance modulus and luminosity distance. Here we correct for Galactic extinction and incorporate the cosmic reddening "k-correction" (Figure 7) into the distance calculation for self-consistency between the measured colors and the inferred luminosity distance.

Independent studies of the K-band luminosity function (Cole et al 2001; Kochanek et al 2001) reveal consistent Schecter Function parameters: $M_\star$ = -24.0 mag with a faint end slope index of about -1.0 (assuming H_0 = 72 km/s/Mpc). We adopt $M_\star$ as the standard candle for 2MASS galaxies. For the faintest 2MASS galaxies, K=14 mag, the implied distance is 400 Mpc or z$\sim$0.1. The actual distribution of Ks-band luminosity for all 2MASS galaxies with known redshifts is shown in Figure 8. The histogram peaks at -24.4 mag or so, implying that 2MASS is sensitive to intrinsically luminous (early-type) galaxies. (Cautionary note: the 2MASS-NED redshift sample comes from a variety of surveys, each with their own biased selection effects – a prime motivation to carry out uniform and minimally-biased redshift survey, such as that of 6dFGS). Approximately 10^5 2MASS galaxies have radial velocity measurements, representing a small fraction ($\sim$ 8%) of the total, but a significant fraction for the largest (nearby) galaxies.

Results – A comparison of the photometric-derived redshifts with radial-velocity redshifts is shown in Figure 9, where we plot the redshift as a function of the Ks-band flux. The photometric redshift accurately predict the mean radial-velocity redshift per mag interval from the brightest (nearby) to the faintest (distant) galaxies. However, note the large scatter in the redshift distribution per mag interval – this is due to galaxies with intrinsically different luminosity (from the brightest ellipticals to the faintest dwarf

© Astronomical Society of Australia 0000



Figure 6. Supergalactic projection of the 2MASS galaxy catalog. Sources are RGB color-coded according to their Ks-band integrated flux: brightest galaxies (Ks $<$ 10) are blue, moderately bright galaxies (10 $<$ Ks $<$ 12.5) are green, and the faintest (Ks $>$ 12.5) are red. The Local Supercluster as defined by de Vaucouleurs et al (1976) extends along the equator crossing through the Virgo galaxy cluster (bright *blue* cluster, left-center of image). The Milky Way is shown with a cyan color scale, crossing the center of the image and outer boundary. The Perseus-Pisces supercluster extends from the center right (northern Galactic hemisphere) down into the ZoA and out the other side to the southern Galactic hemisphere.

galaxies). What this means is that 2MASS photometric redshifts provide the correct answer on average, but for any given galaxy the uncertainty is large, which is particularly severe for the faint end of the distribution. For galaxy groups or clusters the ensemble average redshift should be accurate to $\sim 20\%$.

All in all, photometric redshifts should provide an adequate, if qualitative, representation of the local Universe. Combining the 10^5 radial-velocity redshifts with the 10^6 photometric redshifts, a 3-dimensional construct of the distribution of galaxies comprising the local Universe is created. Figure 1 represents a view into this 3-D construct. Here the 2MASS galaxies are projected onto a Galactic coordinate (equal-area) aitoff grid with 3 arcmin pixels. Distance is denoted with color coding, where blue represents the nearest galaxies (z $<$ 0.01), green at moderate distances (z $\sim$0.03) and red the most distant objects (z $>$ 0.06). To enhance the contrast between the nearest clusters (e.g., Virgo) and the distant universe, the pixel intensity is modified by the integrated flux along the line of sight (analogous to Figure 5). The effect is to brighten the colors of nearby galaxy clusters (e.g., the Virgo cluster appears blue-white in the figure). Finally, the 2MASS point source catalog, representing the Milky Way, is incorporated into the projection. Near the galactic center region (center of image), the confusion noise completely swamps out detection of background extragalactic objects.

The Cosmic Web – Figure 1 beautifully unveils the background extragalactic sky from the obscuring foreground Milky Way. Galaxies are color-coded by their inferred redshift (or distance from the Sun), thereby providing *depth* to the surface distribution of galaxies. Large-scale structures are clearly discerned, from the Local Supercluster (Virgo/Hydra/Centaurus) to the Perseus-Pisces (*P-P*) and Pavo-Indus (*P-I*) superclusters. (Figure 2, a key to the large scale structures). Probing orthogonal to this 3-D surface reveals the redshift distibution along the line of sight; an example is shown in Figure 10 for the Shapley Concentration (discussed below).

As seen in Figure 1, a continuous chain of structures seems to wrap around Hydra/Virgo, up to Coma to the north, through Hercules to the east, down through *P-P* and around to *P-I* to the west, then extending up through the ZoA into the Shapley concentration. This is more easily seen when each redshift layer is shown separately; Figure 11.

The Milky Way and Magellanic Clouds occupy the first (z = 0) redshift layer. The second layer (z $<$ 0.01) is dominated by the Virgo, Fornax and Hydra-Cen superclusters. The third layer (0.01 $<$ z $<$ 0.02) is dominated by the *P-P* supercluster (left side of image) and the *P-I* supercluster extending up into the ZoA terminating as the *Great Attractor* region (notably Abell 3627) disappears behind a wall of Milky Way stars. An intriguing "ring" or chain of galaxies seems to circle/extend from the northern to

© Astronomical Society of Australia 0000



Figure 7. Galaxy near-infrared colors as a function of redshift. Cosmic reddening is the result of shifting of galaxy light from the H-band (1.6 μm) into the Ks-band (2.2 μm).



Figure 8. Luminosity distribution of 92,000 2MASS galaxies as computed from redshift-derived distances. The redshifts were extracted from NED, the bulk of which come from the RC3, UGC, CfA, LCRS, 2dF, 6dFGS and SDSS surveys.



Figure 9. Redshift *vs.* 2MASS Ks-band integrated flux. Photometric redshifts (black curve) are derived assuming $L_\star$ properties for 2MASS galaxies. These are compared to the radial-velocity redshifts (blue points). The error bars represent the scatter in redshift vs. Ks-band integrated flux (i.e., ranging from luminous to sub-luminous galaxies).

the southern Galactic hemisphere (see also Figure 1). It is unknown whether this ring-like structure is real or a projection effect. The fourth layer ($0.02 < z < 0.03$) is characterized by compact galaxy clusters, including Coma (extreme top of image), Abell 3627 in the ZoA, the Ophiuchus cluster directly to the north of the Milky Way center, and clusters associated with *P-P*. The "great wall" of galaxies extends from Coma down toward Bootes and Hercules. The fifth layer ($0.03 < z < 0.04$) is dominated by the Hercules supercluster (top, left of image), the "great wall," Columba supercluster (bottom right of image), and the massive Shapley Concentration is beginning to appear. The sixth layer ($0.04 < z < 0.05$) showcases Abell 3558 of the enormous Shapley Concentration that lurks behind the nearby Hydra-Cen supercluster, and the Sculpter supercluster (bottom of image) makes its first appearance. The seventh layer ($0.05 < z < 0.06$) contains the backside of the Shapley Concentration, while the Sculpter supercluster dominates the southern hemisphere. The eighth and final layer ($z > 0.06$) contains the most distant structures that 2MASS resolves, including the Pisces-Cetus (located behind *P-P*), Bootes (located behind Hercules), Horologium and Corona Borealis galaxy clusters. At these faint flux levels, the photometric redshifts are losing their ability to discern the cosmic web beyond 300 Mpc, smearing and degrading the resolution of the 3-D construct.

This is clearly demonstrated in Figure 10, where we show the redshift distribution for the Shapley Concentration region in a constant declination slice across the equatorial axis. The nearby galaxy clusters (Hydra-Cen) are easily discerned (note the "fingers of god"), as are the rich galaxy clusters of Shapley (center of image), but beyond z ¿ 0.06 the distribution is smoothing out as the uncertainties in the photometric redshifts begin to dominate. But the intricate web of large scale structure extends well beyond this volume limit, as unmistakably demonstrated by the

© Astronomical Society of Australia 0000



Figure 10. Redshift slice of the Shapley Concentration region, extending to the limit of the 2MASS galaxy catalog, z∼0.1 (400 Mpc). The equatorial RA slice is 90° between a declination boundary of $-36.5° < \mathrm{Dec} < -26.5°$. The points are color-coded by redshift (see Figure 1). The "finger of god" radial velocity artifacts trace galaxy clusters, including Abell 3558 (center, yellow). The Hydra-Cen supercluster is foreground (blue fingers) to the Shapley Concentration.

2dF and SDSS galaxy surveys. But with the addition of optical and or mid-infrared photometry from future all sky surveys (e.g., WISE), the photometric redshifts may be greatly improved, allowing reconstruction of the local Universe beyond 300 to 400 Mpc. Moreover, large redshift surveys (e.g., 6dFGS) will provide accurate distance estimates for large regions of the sky, further sharpening our view of the cosmic web.

Final Thoughts – The 2MASS catalog has proven to be quite versatile to the astronomical community: supporting observation and future mission planning, seeding studies of star formation and morphology in nearby galaxies, penetrating the zone of avoidance, providing the base catalog of redshift and Tully-Fisher HI surveys, and so on. But perhaps its most important function is to provide the "big picture" context for analysis and interpretation of data concerning galaxy clusters, large scale structure and the density of matter in the Universe. And so the primary motivation of this work, with the construction of qualitative "road" maps to the local Universe, is to provide a broad framework for studying the physical connection between the local Universe (Milky Way, Local Group, Local Supercluster, "Great Wall", etc) and the distant Universe where galaxies and the cosmic web first formed.



Figure 11. 2MASS galaxy distribution separated by redshift layer. The color scheme, when combined, creates the panorama shown in Figure 1.

5 Acknowledgments

The author would like to thank Joe Mazzarella (NED) and John Huchra (CfA) for kindly providing galaxy redshifts. And a special thanks to Joss Bland-Hawthorn for organizing the LSS workshop (in honor of Brent Tully) that inspired this paper. This publication makes use of data products from 2MASS, which is a joint project of the Univ. of Massachusetts and the Infrared Processing and Analysis Center, funded by the NASA and the NSF. This work was supported in part by the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

REFERENCES

Cole, S. et al 2001, MNRAS, 326, 255.
Courtois, H., Paturel, G., Sousbie, T., & Labini, F. 2004, A&A (in press).
Cutri, R.M., et al 2000, The 2MASS Explanatory Supplement.
De Vaucouleuers, G. , De Vaucouleuers, A., & Corwin, H. 1976, 2nd RGC (Austin: Univ. of Texas Press).
Eisenhardt, P, & Wright, E. 2003, Proceedings of the SPIE, 4850, 1050.
Gardner J.P., Sharples M.R., Frenk C.S., & Carrasco, B.E. 1997, ApJ, 480, L99.
Glazebrook, K., Peakcock, J.A., Collins, C.A., & Miller, L. 1994, MNRAS, 266, 65.
Huchra, J., Davis, M., Latham & Tonry, J., 1983, ApJS 52, 89.
Jarrett, T.H., 2000, PASP, 112, 1008.
Jarrett, T.H., Chester, T., Cutri, R., Schneider, S., Skrutskie, M. & Huchra, J. 2000, AJ, 119, 2498.
Jarrett, T.H., Chester, T., Cutri, R., Schneider, S., Rosenberg, J., & Huchra, J., 2000, AJ, 120, 298.
Jarrett, T. H., Chester, T., Huchra, J., & Schneider, S. 1998, BAAS, 30, 901.
Huchra, J. P., Davis, M., Latham, D., & Tonry, J. 1983, ApJS, 52, 89.
Kochanek, C. S., White, Martin, Huchra, J., Macri, L., Jarrett, T. H., Schneider, S. E., Mader, J. 2003, ApJ, 585, 161.
Kochanek, C.S., et al 2001, ApJ, 560, 566.
Jones, D.H., et al 2004, MNRAS (submitted).
Moore, A.; Gillingham, P.; Saunders, W. 2002, ASP Conference Proceedings, 280, 109.
Skrutskie, M., et al 1997, in The Impact of Large-Scale Near-IR Sky Surveys, ed. F. Garzon et al (Dordrecht: Kluwar), 25.
Tonry, JL, Blakeslee, JP, Ajhar, EA, & Dressler, A. 2000, ApJ, 530, 625.
Tully, R. 1982, ApJ, 257, 389.
Watson, F., et al 2001, ASP Conf. Series vol. 232, pg. 421, Clowes, Adamson and Bromage eds.

© Astronomical Society of Australia 0000